

DR. DARRELL PERKINS

M.B., B.S.(Hons), B.Med.Sc., F.R.A.C.S.
Provider No. 2048605W

AESTHETIC DAY SURGERY

14 KENSINGTON STREET
KOGARAH NSW 2217
TELEPHONE: 9588 5700
FACSIMILE: 9553 9924

DR. DAMIAN MARUCCI

M.B., B.S., B.A., Ph.D., F.R.A.C.S.
Provider No. 206938EX

SUITE 3, LEVEL 4
531 – 533 THE KINGSWAY
MIRANDA NSW 2228
TELEPHONE: 9525 8400

DR. ADRIAN SJARIF

M.B., B.S.(Hons), B.Sc., M.S., F.R.A.C.S.
Provider No. 255103RF

AFTER HOURS
DR. PERKINS: 0408 165 425
DR. MARUCCI: 0413 123 155
DR. SJARIF: 0413 828 414

Post Operative Instructions

SKIN GRAFT CARE

1. **Grafted Area** (*from where the lesion has been removed*)

KEEP DRY

- Leave dressing(s) dry and intact until your review appointment. However, if the outer dressing (gauze and tape) falls off, the remainder dressing underneath (foam rubber) should REMAIN. DO NOT ATTEMPT TO REMOVE THIS DRESSING.
- Keep wound dry and apply Vaseline twice daily until your next appointment

2. a) **Donor Area on Arm or Leg**

- Leave dressings dry and intact until your review appointment.
- Remove the outer bandage in ___ days. Leave the paper tape on the wound. You may get this tape lightly wet in the shower and then pat dry. If the paper tape falls off, replace with new paper tape.
- Remove the outer bandage and Vaseline impregnated gauze in ___ days. You may shower normally. Apply ordinary Vaseline to the wound twice a day.

b) **Donor Area in Front or Behind the Ear or on the Neck**

- Leave dressing(s) dry and intact for 24 hours. You may then shower over the tape.
- If desired, MICROPORE TAPE may be applied directly over the sutures.
- Apply ordinary Vaseline to wound twice daily until your review appointment.

- If you have excessive bleeding, apply DIRECT PRESSURE FOR 10 MINUTES. If necessary, the dressing(s) may be replaced with a similar type. If bleeding persists, ring our rooms or your GP.
- DO NOT shave or apply make-up over the sutures.
- If there is excessive redness, irritation, or any signs of possible infection ring our rooms or your GP.
- If your wound is on your arm or leg, keep that limb elevated for the first 2 days, and whenever possible until the wound is completely healed.
- If required, pain medication may be taken (not Aspirin).

Follow up appointment:

at: Kogarah Rooms Miranda Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

