

DR. DARRELL PERKINS

M.B., B.S.(Hons), B.Med.Sc., F.R.A.C.S.
Provider No. 2048605W

AESTHETIC DAY SURGERY

14 KENSINGTON STREET
KOGARAH NSW 2217
TELEPHONE: 9588 5700
FACSIMILE: 9553 9924

DR. DAMIAN MARUCCI

M.B., B.S., B.A., Ph.D., F.R.A.C.S.
Provider No. 206938EX

SUITE 3, LEVEL 4
531 – 533 THE KINGSWAY
MIRANDA NSW 2228
TELEPHONE: 9525 8400

DR. ADRIAN SJARIF

M.B., B.S.(Hons), B.Sc., M.S., F.R.A.C.S.
Provider No. 255103RF

AFTER HOURS
DR. PERKINS: 0408 165 425
DR. MARUCCI: 0413 123 155
DR. SJARIF: 0413 828 414

Post Operative Instructions

FACELIFT

1. Take all medications as directed
2. Sleep on 2 or 3 pillows to keep the head elevated. This will reduce swelling
3. Avoid bending down. Keep your head elevated at all times
4. Apply ice packs to the face as much as possible for the first 48 hours, eg. for 20 minutes every 1 – 2 hours
5. REST ONLY. Do no exercise. Keep cool
6. Expect some swelling of the face. A small amount of blood ooze through the bandage can also be expected

Further instructions will be given when we remove the dressings in two days. Please bring a scarf with you to this appointment

EYELID REDUCTION

7. Use cold saline and cold water (boil first then chill) packs frequently to reduce swelling and soothe discomfort
8. Clean the eyes with a moist cotton bud 3 times a day
9. **If there is severe pain or visual impairment contact us immediately** – this would be an extremely unlikely occurrence

For lubrication of the eyes purchase:

(Remove soft contact lenses before using)

Day: “Refresh” or “Murine Revital Eyes” or similar eye lubricant

Night: “Viscotears Liquid Gel”

Please take care when applying lubricant to your eyes not to pull on the upper or lower eyelid wounds

Follow up appointment:

at: Kogarah Rooms Miranda Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

Please see the reverse for possible adverse reactions and drug interactions →



Possible adverse reactions and drug interactions:

Medication	Possible Adverse Reaction
Refresh Murine Revital Eyes	Stinging and burning
Viscotears Liquid Gel	Mild transient burning, sticky eyelids, blurred vision

If you encounter any adverse reaction/s to any of your medications, cease the medication and contact your Surgeon (refer to the Post Operative Instructions sheet) for further advice.