

DR. DARRELL PERKINS

M.B., B.S.(Hons), B.Med.Sc., F.R.A.C.S.
Provider No. 2048605W

DR. DAMIAN MARUCCI

M.B., B.S., B.A., Ph.D., F.R.A.C.S.
Provider No. 206938EX

DR. ADRIAN SJARIF

M.B., B.S.(Hons), B.Sc., M.S., F.R.A.C.S.
Provider No. 255103RF

AESTHETIC DAY SURGERY

14 KENSINGTON STREET
KOGARAH NSW 2217
TELEPHONE: 9588 5700
FACSIMILE: 9553 9924

SUITE 3, LEVEL 4
531 – 533 THE KINGSWAY
MIRANDA NSW 2228
TELEPHONE: 9525 8400

AFTER HOURS
DR. PERKINS: 0408 165 425
DR. MARUCCI: 0413 123 155
DR. SJARIF: 0413 828 414

Post Operative Instructions

LABIOPLASTY

1. For the first two days after surgery take 2 Panadol every six hours for pain relief. **DO NOT TAKE ASPIRIN**
2. Take frequent warm showers to reduce discomfort
3. Use a hairdryer on a low setting to dry the stitches after bathing, and try to keep the area as dry as possible
4. Change pads frequently

Follow up appointment:

at: Kogarah Rooms Miranda Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

Please see the reverse for possible adverse reactions and drug interactions →



Medication	Possible Adverse Reaction
Panadol (Paracetamol)	(Rare) Stomach upset, nausea, rash [see below]

NB: Paracetamol may be used to treat pain, or temperatures over 38°C. It is not to be used for mild fever, gastroenteritis or as a sedative.

Oral Paracetamol dosage recommendations for adults and children after puberty:
Use Paracetamol 4 to 6 hourly, with a maximum of 6 tablets daily, and for no more than 48 – 36 hours for patients of normal or average build. If pain persists seek medical advice before continuing.

All patients taking Paracetamol for extended periods should be aware of the Paracetamol content in over the counter and prescription products that they are taking as there could be an increased risk of liver problems.

If you encounter any adverse reaction/s to any of your medications, cease the medication and contact your Surgeon (refer to the Post Operative Instructions sheet) for further advice.