

**DR. DARRELL PERKINS**

MBBS(Hons), BMedSc, FRACS  
Provider No. 2048605W

14 KENSINGTON STREET  
KOGARAH NSW 2217  
TELEPHONE: 9588 5700

**DR. DAMIAN MARUCCI**

MBBS, BA, PhD, FRACS  
Provider No. 206938EX

531 – 533 THE KINGSWAY  
MIRANDA NSW 2228  
TELEPHONE: 9525 8400

**DR. ADRIAN SJARIF**

MBBS(Hons), BSc, MS, FRACS  
Provider No. 255103RF

372 CROWN STREET  
WOLLONGONG NSW 2500  
TELEPHONE: 4226 6111

AFTER HOURS  
DR. PERKINS: 0408 165 425  
DR. MARUCCI: 0413 123 155  
DR. SJARIF: 0413 828 414

Post Operative Instructions

**FACELIFT**

1. Take all medications as directed
2. Sleep on 2 or 3 pillows to keep the head elevated. This will reduce swelling
3. Avoid bending down. Keep your head elevated at all times
4. Apply ice packs to the face as much as possible for the first 48 hours, e.g. for 20 minutes every 1 – 2 hours
5. REST ONLY. Do no exercise. Keep cool
6. Expect some swelling of the face. A small amount of blood ooze through the bandage can also be expected.

Further instructions will be given when we remove the dressings in two days. Please bring a scarf with you to this appointment

**EYELID REDUCTION**

7. Use cold saline and cold water (boil first then chill) packs frequently to reduce swelling and soothe discomfort
8. Clean the eyes with a moist cotton bud 3 times a day
9. **If there is severe pain or visual impairment contact us immediately** – this would be an extremely unlikely occurrence.

For lubrication of the eyes (do not purchase until advised by surgeon):

Remove soft contact lenses before using

**Day:** “Refresh” or “Murine Revital Eyes” or similar eye lubricant

**Night:** “Viscotears Liquid Gel”

**Follow up appointment:** .....

at:  Kogarah Rooms  Miranda Rooms  Wollongong Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

**Please see the reverse for possible adverse reactions and drug interactions →**

**Possible adverse reactions and drug interactions:**

<b>Medication</b>	<b>Possible Adverse Reaction</b>
Refresh Murine Revital Eyes	Stinging and burning
Viscotears Liquid Gel	Mild transient burning, sticky eyelids, blurred vision

If you encounter any adverse reaction/s to any of your medications, cease the medication and contact your Surgeon (refer to the Post Operative Instructions sheet) for further advice.

