

**DR. DARRELL PERKINS**

M.B., B.S.(Hons), B.Med.Sc., F.R.A.C.S.  
Provider No. 2048605W

**DR. DAMIAN MARUCCI**

M.B., B.S., B.A., Ph.D., F.R.A.C.S.  
Provider No. 206938EX

**DR. ADRIAN SJARIF**

M.B., B.S.(Hons), B.Sc., M.S., F.R.A.C.S.  
Provider No. 255103RF

**AESTHETIC DAY SURGERY**

14 KENSINGTON STREET  
KOGARAH NSW 2217  
TELEPHONE: 9588 5700  
FACSIMILE: 9553 9924

SUITE 3, LEVEL 4  
531 – 533 THE KINGSWAY  
MIRANDA NSW 2228  
TELEPHONE: 9525 8400

AFTER HOURS  
DR. PERKINS: 0408 165 425  
DR. MARUCCI: 0413 123 155  
DR. SJARIF: 0413 828 414

Post Operative Instructions

**CORRECTION OF PROMINENT EARS**

1. Keep crepe bandage dry and intact. If this comes off before the first visit it will need to be re-done
2. Take pain relief as prescribed
3. For severe itching take Phenergan, dose as prescribed by the Pharmacist

Follow up appointment: .....

at:  Kogarah Rooms  Miranda Rooms

If you have any questions or concerns, please contact us on one of the numbers above.

**Please see the reverse for possible adverse reactions and drug interactions →**



<b>Medication</b>	<b>Possible Adverse Reaction</b>
Panadol (Paracetamol)	(Rare) Stomach upset, nausea, rash [see below]
Phenergan	Sedation, excitation, convulsions, nausea and vomiting, dry mouth, palpitations, vision changes

NB: Paracetamol may be used to treat pain, or temperatures over 38°C. It is not to be used for mild fever, gastroenteritis or as a sedative.

*Oral Paracetamol dosage recommendations for children up to puberty:*

Follow dosage instructions on the packet. In all cases, Paracetamol should not be administered to children up to puberty for more than 48 hours without medical review.

*Oral Paracetamol dosage recommendations for adults and children after puberty:*

Use Paracetamol 4 to 6 hourly, with a maximum of 6 tablets daily, and for no more than 48 – 36 hours for patients of normal or average build. If pain persists seek medical advice before continuing.

All patients taking Paracetamol for extended periods should be aware of the Paracetamol content in over the counter and prescription products that they are taking as there could be an increased risk of liver problems.

If you encounter any adverse reaction/s to any of your medications, cease the medication and contact your Surgeon (refer to the Post Operative Instructions sheet) for further advice.